

HEALTH MONITORING REPORT (November 2015)

Species: MICE

Agents	Area	A11	A12	A14
	Strains & Stocks	CAnN.Cg-Foxn1 <sup>nu</sup> /CrjCrlj <sup>3</sup> Crlj:CD1-Foxn1 <sup>nu</sup> <sup>3</sup>	CAnN.Cg-Foxn1 <sup>nu</sup> /CrjCrlj <sup>3</sup>	JAX <sup>®</sup> Mice Strain SJL/J JAX <sup>®</sup> Mice Strain CBA/J NC/NgaTndCrlj Crl:CD1(ICR)
	Test period Technique	November 9 - 21	November 9 - 21	November 9 - 21
<b>VIRUSES</b>				
Sendai Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Pneumonia Virus of Mice <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Mouse Hepatitis Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Minute Virus of Mice <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Mouse Parvovirus (1 or 2) <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Murine Norovirus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Theiler's Murine Encephalomyelitis Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Reovirus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Epizootic Diarrhea of Infant Mice Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Lymphocytic Choriomeningitis Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Ectromelia Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Adenovirus (1 or 2) <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Cytomegalovirus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Pneumonitis Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Polyoma Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Hantaan Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Thymic Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<b>BACTERIA</b>				
<i>Bordetella bronchiseptica</i> (m)	Culture	0/12	0/12	0/8
<i>Citrobacter rodentium</i> (m)	Culture	0/8	0/8	0/8
<i>Corynebacterium kutscheri</i> (m)	Culture	0/12	0/12	0/8
<i>Pasteurella pneumotropica</i> (m)	Culture	0/12	0/12	0/8
<i>Pseudomonas aeruginosa</i> <sup>2</sup> (m)	Culture	0/12	0/12	0/8
<i>Salmonella</i> spp. (m)	Culture	0/12	0/12	0/8
<i>Streptococcus pneumoniae</i> (m)	Culture	0/12	0/12	0/8
<i>Staphylococcus aureus</i> <sup>2</sup> (m)	Culture	0/12	0/12	-
<i>Helicobacter hepaticus</i> (q)	PCR	0/12	0/12	0/8
<i>Helicobacter bilis</i> (q)	PCR	0/12	0/12	0/8
<i>Helicobacter</i> spp. (q)	PCR	0/12	0/12	0/8
<i>Streptobacillus moniliformis</i> (q)	PCR	0/12	0/12	0/8
<i>Mycoplasma pulmonis</i> <sup>1</sup> (m)	PCR/MFIA	0/12	0/12	0/8
<i>Clostridium piliforme</i> (Tyzzer's Disease) <sup>1</sup> (m)	Lesions/MFIA	0/12	0/12	0/12
CAR bacillus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<b>PARASITES AND FUNGI</b>				
Ectoparasites (m)	Direct Examination	0/12	0/12	0/8
Pinworms (m)	Direct Examination	0/12	0/12	0/12
Gastrointestinal protozoa (m)	Direct Examination	0/8	0/8	0/8
<i>Encephalitozoon cuniculi</i> <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<i>Pneumocystis</i> spp. <sup>2</sup> (q)	PCR	0/4	0/4	-

Monitoring frequency: (m) monthly, (q) quarterly

Remarks: 1: Serology on immunocompetent animals from each colony

2: Not considered as specific pathogens for immunocompetent strains

3: Monitoring on both *nu/nu* and *nu/+* mice

Note: JAX<sup>®</sup> is a registered trademark of The Jackson Laboratory. All rights reserved.

Tested by Charles River Laboratories Japan, Inc.

日本チャールス・リバー株式会社  
代表取締役社長 小林 功

HEALTH MONITORING REPORT (November 2015)

Species: MICE

Agents	Area	A23	A24	A25
	Strains & Stocks	BALB/cAnNCrIrlj DBA/2NCrIrlj CD2F1/CrIrlj <sup>3</sup>	CAnN.Cg-Foxn1 <sup>nu</sup> /CrIrlj <sup>4</sup>	JAX® Mice Strain C57BL/6J
	Test period Technique	November 2 - 21	November 9 - 21	November 9 - 25
<b>VIRUSES</b>				
Sendai Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Pneumonia Virus of Mice <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Mouse Hepatitis Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Minute Virus of Mice <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Mouse Parvovirus (1 or 2) <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Murine Norovirus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Theiler's Murine Encephalomyelitis Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Reovirus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Epizootic Diarrhea of Infant Mice Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Lymphocytic Choriomeningitis Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Ectromelia Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Adenovirus (1 or 2) <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Cytomegalovirus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Pneumonitis Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Polyoma Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Hantaan Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Thymic Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<b>BACTERIA</b>				
<i>Bordetella bronchiseptica</i> (m)	Culture	0/8	0/12	0/8
<i>Citrobacter rodentium</i> (m)	Culture	0/8	0/8	0/8
<i>Corynebacterium kutscheri</i> (m)	Culture	0/8	0/12	0/8
<i>Pasteurella pneumotropica</i> (m)	Culture	0/8	0/12	0/8
<i>Pseudomonas aeruginosa</i> <sup>2</sup> (m)	Culture	0/8	0/12	0/8
<i>Salmonella</i> spp. (m)	Culture	0/8	0/12	0/8
<i>Streptococcus pneumoniae</i> (m)	Culture	0/8	0/12	0/8
<i>Staphylococcus aureus</i> <sup>2</sup> (m)	Culture	-	0/12	-
<i>Helicobacter hepaticus</i> (q)	PCR	0/8	0/12	0/8
<i>Helicobacter bilis</i> (q)	PCR	0/8	0/12	0/8
<i>Helicobacter</i> spp. (q)	PCR	0/8	0/12	0/8
<i>Streptobacillus moniliformis</i> (q)	PCR	0/8	0/12	0/8
<i>Mycoplasma pulmonis</i> <sup>1</sup> (m)	PCR/MFIA	0/8	0/12	0/8
<i>Clostridium piliforme</i> (Tyzzer's Disease) <sup>1</sup> (m)	Lesions/MFIA	0/12	0/12	0/12
CAR bacillus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<b>PARASITES AND FUNGI</b>				
Ectoparasites (m)	Direct Examination	0/8	0/12	0/8
Pinworms (m)	Direct Examination	0/12	0/12	0/12
Gastrointestinal protozoa (m)	Direct Examination	0/8	0/8	0/8
<i>Encephalitozoon cuniculi</i> <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<i>Pneumocystis</i> spp. <sup>2</sup> (q)	PCR	-	0/4	-

Tested by Charles River Laboratories Japan, Inc.

Monitoring frequency: (m) monthly, (q) quarterly

Remarks: 1: Serology on immunocompetent animals from each colony

2: Not considered as specific pathogens for immunocompetent strains

3: Not tested strains or stocks; monitoring on parents or other strains in the area

4: Monitoring on both nu/nu and nu/+ mice

Note: JAX® is a registered trademark of The Jackson Laboratory. All rights reserved.

日本チャールス・リバー株式会社  
代表取締役社長 小林 功

HEALTH MONITORING REPORT (November 2015)

Species: RATS

Agents	Area	A21	A22
	Strains & Stocks	F344/DuCrjCrj ZDF- <i>Lepr<sup>fa</sup></i> /CrjCrj Crj:ZUC- <i>Lepr<sup>fa</sup></i>	Crj:CD(SD)
	Test period Technique	November 2 - 21	November 2 - 21
<b>VIRUSES</b>			
Sendai Virus <sup>1</sup> (m)	MFIA	0/8	0/8
Pneumonia Virus of Mice <sup>1</sup> (m)	MFIA	0/8	0/8
Sialodacryoadenitis Virus <sup>1</sup> (m)	MFIA	0/8	0/8
Rat Minute Virus <sup>1</sup> (m)	MFIA	0/8	0/8
Rat Parvovirus <sup>1</sup> (m)	MFIA	0/8	0/8
Toolan's H-1 Virus <sup>1</sup> (m)	MFIA	0/8	0/8
Kilham Rat Virus <sup>1</sup> (m)	MFIA	0/8	0/8
Reovirus <sup>1</sup> (m)	MFIA	0/8	0/8
Rat Theilovirus <sup>1</sup> (m)	MFIA	0/8	0/8
Lymphocytic Choriomeningitis Virus <sup>1</sup> (q)	MFIA	0/8	0/8
Mouse Adenovirus (1 or 2) <sup>1</sup> (q)	MFIA	0/8	0/8
Hantaan Virus <sup>1</sup> (q)	MFIA	0/8	0/8
<b>BACTERIA</b>			
<i>Bordetella bronchiseptica</i> (m)	Culture	0/8	0/8
<i>Corynebacterium kutscheri</i> (m)	Culture	0/8	0/8
<i>Pasteurella pneumotropica</i> (m)	Culture	0/8	0/8
<i>Pseudomonas aeruginosa</i> <sup>2</sup> (m)	Culture	0/8	0/8
<i>Salmonella</i> spp. (m)	Culture	0/8	0/8
<i>Streptococcus pneumoniae</i> (m)	Culture	0/8	0/8
<i>Helicobacter hepaticus</i> (q)	PCR	0/8	0/8
<i>Helicobacter bilis</i> (q)	PCR	0/8	0/8
<i>Helicobacter</i> spp. (q)	PCR	0/8	0/8
<i>Streptobacillus moniliformis</i> (q)	PCR	0/8	0/8
<i>Mycoplasma pulmonis</i> <sup>1</sup> (m)	PCR/MFIA	0/8	0/8
<i>Clostridium piliforme</i> (Tyzzer's Disease) <sup>1</sup> (m)	Lesions/MFIA	0/12	0/12
CAR bacillus <sup>1</sup> (q)	MFIA	0/8	0/8
<b>PARASITES AND FUNGI</b>			
Ectoparasites (m)	Direct Examination	0/8	0/8
Pinworms (m)	Direct Examination	0/12	0/12
Gastrointestinal protozoa (m)	Direct Examination	0/8	0/8
<i>Encephalitozoon cuniculi</i> <sup>1</sup> (q)	MFIA	0/8	0/8
<i>Pneumocystis</i> spp. <sup>1,3</sup> (m)	MFIA	0/8	0/8

Tested by Charles River Laboratories Japan, Inc.

Monitoring frequency: (m) monthly, (q) quarterly

Remarks: 1: Serology on immunocompetent animals from each colony  
2: Not considered as specific pathogens for immunocompetent strains  
3: Not included in the Specific Pathogen Free (SPF) lists

Agents	Area	H12	H22	H23
	Strains & Stocks	JAX <sup>®</sup> Mice Strain C57BL/6J JAX <sup>®</sup> Mice Strain B6.Cg-Lep <sup>ob</sup> /J JAX <sup>®</sup> Mice Strain BKS.Cg-Dock7 <sup>m</sup> +/+ Lep <sup>ob</sup> /J	CrI:CD1(ICR) C57BL/6NCrI <sup>3</sup> C3H/HeNCrI DBA/2NCrI B6N-Ty <sup>c-Brd</sup> /BrdCrCrI B6D2F1/CrI <sup>3</sup> B6C3F1/CrI <sup>3</sup>	BALB/cAnNCrI DBA/1JNCrI
	Test period Technique	November 4 - 21	November 4 - 21	November 4 - 21
<b>VIRUSES</b>				
Sendai Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Pneumonia Virus of Mice <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Mouse Hepatitis Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Minute Virus of Mice <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Mouse Parvovirus (1 or 2) <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Murine Norovirus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Theiler's Murine Encephalomyelitis Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Reovirus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Epizootic Diarrhea of Infant Mice Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Lymphocytic Choriomeningitis Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Ectromelia Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Adenovirus (1 or 2) <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Cytomegalovirus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Pneumonitis Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Polyoma Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Hantaan Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Thymic Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<b>BACTERIA</b>				
<i>Bordetella bronchiseptica</i> (m)	Culture	0/8	0/8	0/8
<i>Citrobacter rodentium</i> (m)	Culture	0/8	0/8	0/8
<i>Corynebacterium kutscheri</i> (m)	Culture	0/8	0/8	0/8
<i>Pasteurella pneumotropica</i> (m)	Culture	0/8	0/8	0/8
<i>Pseudomonas aeruginosa</i> <sup>2</sup> (m)	Culture	0/8	0/8	0/8
<i>Salmonella</i> spp. (m)	Culture	0/8	0/8	0/8
<i>Streptococcus pneumoniae</i> (m)	Culture	0/8	0/8	0/8
<i>Helicobacter hepaticus</i> (q)	PCR	0/8	0/8	0/8
<i>Helicobacter bilis</i> (q)	PCR	0/8	0/8	0/8
<i>Helicobacter</i> spp. (q)	PCR	0/8	0/8	0/8
<i>Streptobacillus moniliformis</i> (q)	PCR	0/8	0/8	0/8
<i>Mycoplasma pulmonis</i> <sup>1</sup> (m)	PCR/MFIA	0/8	0/8	0/8
<i>Clostridium piliforme</i> (Tyzzer's Disease) <sup>1</sup> (m)	Lesions/MFIA	0/12	0/12	0/12
CAR bacillus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<b>PARASITES AND FUNGI</b>				
Ectoparasites (m)	Direct Examination	0/8	0/8	0/8
Pinworms (m)	Direct Examination	0/12	0/12	0/12
Gastrointestinal protozoa (m)	Direct Examination	0/8	0/8	0/8
<i>Encephalitozoon cuniculi</i> <sup>1</sup> (q)	MFIA	0/8	0/8	0/8

Tested by Charles River Laboratories Japan, Inc.

Monitoring frequency: (m) monthly, (q) quarterly

Remarks: 1: Serology on immunocompetent animals from each colony

2: Not considered as specific pathogens for immunocompetent strains

3: Not tested strains or stocks; monitoring on parents or other strains in the area

Note: JAX<sup>®</sup> is a registered trademark of The Jackson Laboratory. All rights reserved.

日本チャールス・リバー株式会社  
代表取締役社長 小林 功

HEALTH MONITORING REPORT (November 2015)

Species: RATS

Agents	Area	H11	H13	H21
	Strains & Stocks	CrI:WI(Han) SHR/NCrI:CrIj WKY/NCrI:CrIj CrIj:WI	LEW/CrI:CrIj BN/CrI:CrIj PCK/CrIjCrI-Pkhd1 <sup>Pck</sup> /CrI:CrIj CrIj:LE	CrI:CD(SD)
	Test period Technique	November 11 - 25	November 11 - 21	November 11 - 21
<b>VIRUSES</b>				
Sendai Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Pneumonia Virus of Mice <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Sialodacryoadenitis Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Rat Minute Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Rat Parvovirus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Toolan's H-1 Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Kilham Rat Virus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Reovirus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Rat Theilovirus <sup>1</sup> (m)	MFIA	0/8	0/8	0/8
Lymphocytic Choriomeningitis Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Mouse Adenovirus (1 or 2) <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
Hantaan Virus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<b>BACTERIA</b>				
<i>Bordetella bronchiseptica</i> (m)	Culture	0/8	0/8	0/8
<i>Corynebacterium kutscheri</i> (m)	Culture	0/8	0/8	0/8
<i>Pasteurella pneumotropica</i> (m)	Culture	0/8	0/8	0/8
<i>Pseudomonas aeruginosa</i> <sup>2</sup> (m)	Culture	0/8	0/8	0/8
<i>Salmonella</i> spp. (m)	Culture	0/8	0/8	0/8
<i>Streptococcus pneumoniae</i> (m)	Culture	0/8	0/8	0/8
<i>Helicobacter hepaticus</i> (q)	PCR	0/8	0/8	0/8
<i>Helicobacter bilis</i> (q)	PCR	0/8	0/8	0/8
<i>Helicobacter</i> spp. (q)	PCR	0/8	0/8	0/8
<i>Streptobacillus moniliformis</i> (q)	PCR	0/8	0/8	0/8
<i>Mycoplasma pulmonis</i> <sup>1</sup> (m)	PCR/MFIA	0/8	0/8	0/8
<i>Clostridium piliforme</i> (Tyzzer's Disease) <sup>1</sup> (m)	Lesions/MFIA	0/12	0/12	0/12
CAR bacillus <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<b>PARASITES AND FUNGI</b>				
Ectoparasites (m)	Direct Examination	0/8	0/8	0/8
Pinworms (m)	Direct Examination	0/12	0/12	0/12
Gastrointestinal protozoa (m)	Direct Examination	0/8	0/8	0/8
<i>Encephalitozoon cuniculi</i> <sup>1</sup> (q)	MFIA	0/8	0/8	0/8
<i>Pneumocystis</i> spp. <sup>1,3</sup> (m)	MFIA	0/8	0/8	0/8

Tested by Charles River Laboratories Japan, Inc.

Monitoring frequency: (m) monthly, (q) quarterly

Remarks: 1: Serology on immunocompetent animals from each colony  
2: Not considered as specific pathogens for immunocompetent strains  
3: Not included in the Specific Pathogen Free (SPF) lists

日本チャールス・リバー株式会社  
代表取締役社長 小林 功

HEALTH MONITORING REPORT (November 2015)

Species: MICE

Agents	Area	T21 Isolator	T21 Isolator	T21 Isolator
	Strains & Stocks	JAX <sup>®</sup> Mice Strain NOD.Cg-Prkdc <sup>scid</sup> Il2rg <sup>tm1Wjl</sup> /SzJ C3H/HeNCrIcrIj (Sentinels)	CrIj:SHO-Prkdc <sup>scid</sup> H <sup>hr</sup> C3H/HeNCrIcrIj (Sentinels)	CB17.Cg-Prkdc <sup>scid</sup> Lysf <sup>bg-J</sup> /CrIcrIj C3H/HeNCrIcrIj (Sentinels)
	Test period Technique	November 4 - 21	November 4 - 21	November 4 - 21
<b>VIRUSES</b>				
Sendai Virus <sup>1</sup> (q)	MFIA	0/7	0/2	0/3
Pneumonia Virus of Mice <sup>1</sup> (q)	MFIA	0/7	0/2	0/3
Mouse Hepatitis Virus <sup>1</sup> (q)	MFIA	0/7	0/2	0/3
Minute Virus of Mice <sup>1</sup> (q)	MFIA	0/7	0/2	0/3
Mouse Parvovirus (1 or 2) <sup>1</sup> (q)	MFIA	0/7	0/2	0/3
Murine Norovirus <sup>1</sup> (q)	MFIA	0/7	0/2	0/3
Theiler's Murine Encephalomyelitis Virus <sup>1</sup> (q)	MFIA	0/7	0/2	0/3
Reovirus <sup>1</sup> (q)	MFIA	0/7	0/2	0/3
Epizootic Diarrhea of Infant Mice Virus <sup>1</sup> (q)	MFIA	0/7	0/2	0/3
<b>BACTERIA</b>				
<i>Bordetella bronchiseptica</i> (q)	Culture	0/21	0/6	0/9
<i>Citrobacter rodentium</i> (q)	Culture	0/14	0/4	0/6
<i>Corynebacterium kutscheri</i> (q)	Culture	0/21	0/6	0/9
<i>Pasteurella pneumotropica</i> (q)	Culture	0/21	0/6	0/9
<i>Pseudomonas aeruginosa</i> (q)	Culture	0/21	0/6	0/9
<i>Salmonella</i> spp. (q)	Culture	0/21	0/6	0/9
<i>Staphylococcus aureus</i> (q)	Culture	0/21	0/6	0/9
<i>Streptococcus pneumoniae</i> (q)	Culture	0/21	0/6	0/9
<i>Mycoplasma pulmonis</i> <sup>1</sup> (q)	MFIA	0/7	0/2	0/3
<i>Clostridium piliforme</i> (Tyzzer's Disease) <sup>1</sup> (q)	Lesions/MFIA	0/21	0/6	0/9
<b>PARASITES AND FUNGI</b>				
Ectoparasites (q)	Direct Examination	0/21	0/6	0/9
Pinworms (q)	Direct Examination	0/21	0/6	0/9
Gastrointestinal protozoa (q)	Direct Examination	0/14	0/4	0/6
<i>Pneumocystis</i> spp. (q)	PCR	0/7	0/2	0/3

Tested by Charles River Laboratories Japan, Inc.

Monitoring frequency of animals: (q) quarterly, (a) annually

Remarks: 1: Serology on immunocompetent animals from each colony

Note: JAX<sup>®</sup> is a registered trademark of The Jackson Laboratory. All rights reserved.

日本チャールス・リバー株式会社  
代表取締役社長 小林 功

HEALTH MONITORING REPORT (November 2015)

Printed date: November 28, 2015

Location: Tsukuba Breeding Center

Species: MICE

Agents	Area	T22 Isolator
	Strains & Stocks	CB17/Icr-Prkdc <sup>scid</sup> /CrIcrlj C3H/HeNcrIcrlj (Sentinels)
	Test period Technique	November 2 - 21
<b>VIRUSES</b>		
Sendai Virus <sup>1</sup> (q)	MFIA	0/10
Pneumonia Virus of Mice <sup>1</sup> (q)	MFIA	0/10
Mouse Hepatitis Virus <sup>1</sup> (q)	MFIA	0/10
Minute Virus of Mice <sup>1</sup> (q)	MFIA	0/10
Mouse Parvovirus (1 or 2) <sup>1</sup> (q)	MFIA	0/10
Murine Norovirus <sup>1</sup> (q)	MFIA	0/10
Theiler's Murine Encephalomyelitis Virus <sup>1</sup> (q)	MFIA	0/10
Reovirus <sup>1</sup> (q)	MFIA	0/10
Epizootic Diarrhea of Infant Mice Virus <sup>1</sup> (q)	MFIA	0/10
<b>BACTERIA</b>		
<i>Bordetella bronchiseptica</i> (q)	Culture	0/30
<i>Citrobacter rodentium</i> (q)	Culture	0/20
<i>Corynebacterium kutscheri</i> (q)	Culture	0/30
<i>Pasteurella pneumotropica</i> (q)	Culture	0/30
<i>Pseudomonas aeruginosa</i> (q)	Culture	0/30
<i>Salmonella</i> spp. (q)	Culture	0/30
<i>Staphylococcus aureus</i> (q)	Culture	0/30
<i>Streptococcus pneumoniae</i> (q)	Culture	0/30
<i>Mycoplasma pulmonis</i> <sup>1</sup> (q)	MFIA	0/10
<i>Clostridium piliforme</i> (Tyzzer's Disease) <sup>1</sup> (q)	Lesions/MFIA	0/30
<b>PARASITES AND FUNGI</b>		
Ectoparasites (q)	Direct Examination	0/30
Pinworms (q)	Direct Examination	0/30
Gastrointestinal protozoa (q)	Direct Examination	0/20
<i>Pneumocystis</i> spp. (q)	PCR	0/10

Tested by Charles River Laboratories Japan, Inc.

Monitoring frequency of animals: (q) quarterly, (a) annually

日本チャールス・リバー株式会社  
代表取締役社長 小林 功

Remarks: 1: Serology on immunocompetent animals from each colony

HEALTH MONITORING REPORT (November 2015)

Printed date: November 28, 2015

Location: Tsukuba Breeding Center

Species: MICE

Agents	Area	T23 Isolator
	Strains & Stocks	JAX <sup>®</sup> Mice Strain NOD.CB17-Prkdc <sup>scid</sup> /J C3H/HeNCrCrj (Sentinels)
	Test period Technique	November 9 - 25
<b>VIRUSES</b>		
Sendai Virus <sup>1</sup> (q)	MFIA	0/7
Pneumonia Virus of Mice <sup>1</sup> (q)	MFIA	0/7
Mouse Hepatitis Virus <sup>1</sup> (q)	MFIA	0/7
Minute Virus of Mice <sup>1</sup> (q)	MFIA	0/7
Mouse Parvovirus (1 or 2) <sup>1</sup> (q)	MFIA	0/7
Murine Norovirus <sup>1</sup> (q)	MFIA	0/7
Theiler's Murine Encephalomyelitis Virus <sup>1</sup> (q)	MFIA	0/7
Reovirus <sup>1</sup> (q)	MFIA	0/7
Epizootic Diarrhea of Infant Mice Virus <sup>1</sup> (q)	MFIA	0/7
<b>BACTERIA</b>		
<i>Bordetella bronchiseptica</i> (q)	Culture	0/21
<i>Citrobacter rodentium</i> (q)	Culture	0/14
<i>Corynebacterium kutscheri</i> (q)	Culture	0/21
<i>Pasteurella pneumotropica</i> (q)	Culture	0/21
<i>Pseudomonas aeruginosa</i> (q)	Culture	0/21
<i>Salmonella</i> spp. (q)	Culture	0/21
<i>Staphylococcus aureus</i> (q)	Culture	0/21
<i>Streptococcus pneumoniae</i> (q)	Culture	0/21
<i>Mycoplasma pulmonis</i> <sup>1</sup> (q)	MFIA	0/7
<i>Clostridium piliforme</i> (Tyzzer's Disease) <sup>1</sup> (q)	Lesions/MFIA	0/21
<b>PARASITES AND FUNGI</b>		
Ectoparasites (q)	Direct Examination	0/21
Pinworms (q)	Direct Examination	0/21
Gastrointestinal protozoa (q)	Direct Examination	0/14
<i>Pneumocystis</i> spp. (q)	PCR	0/7

Tested by Charles River Laboratories Japan, Inc.

Monitoring frequency of animals: (q) quarterly, (a) annually

Remarks: 1: Serology on immunocompetent animals from each colony

Note: JAX<sup>®</sup> is a registered trademark of The Jackson Laboratory. All rights reserved.

日本チャールス・リバー株式会社  
代表取締役社長 小林 功



HEALTH MONITORING REPORT (November 2015)

Printed date: November 28, 2015  
Location: Tsukuba Breeding Center

Species: MICE (Fecal and Water Samples)

Agents	Area	T21 Isolator	T21 Isolator	T21 Isolator
	Strains & Stocks	JAX <sup>®</sup> Mice Strain NOD.Cg-Prkdc <sup>scid</sup> Il2rg <sup>tm1Wjl</sup> /SzJ C3H/HeNCrCrIj (Sentinels)	CrIj:SHO-Prkdc <sup>scid</sup> Hr <sup>hr</sup> C3H/HeNCrCrIj (Sentinels)	CB17.Cg-Prkdc <sup>scid</sup> Lyst <sup>bg-J</sup> /CrIj C3H/HeNCrCrIj (Sentinels)
	Test period Technique	November 16 - 26	November 16 - 26	November 16 - 26
<b>BACTERIA</b>				
<i>Bordetella bronchiseptica</i>	Culture	0/15	0/5	0/5
<i>Citrobacter rodentium</i>	Culture	0/15	0/5	0/5
<i>Corynebacterium kutscheri</i>	Culture	0/15	0/5	0/5
<i>Pasteurella pneumotropica</i>	Culture	0/15	0/5	0/5
<i>Pseudomonas aeruginosa</i>	Culture	0/15	0/5	0/5
<i>Salmonella</i> spp.	Culture	0/15	0/5	0/5
<i>Staphylococcus aureus</i>	Culture	0/15	0/5	0/5
<i>Streptococcus pneumoniae</i>	Culture	0/15	0/5	0/5

Tested by Charles River Laboratories Japan, Inc.

Monitoring frequency: Fecal and water samples of isolator-maintained animals are tested monthly.

HEALTH MONITORING REPORT (November 2015)

Printed date: November 28, 2015  
Location: Tsukuba Breeding Center

Species: MICE (Fecal and Water Samples)

Agents	Area	T22 Isolator
	Strains & Stocks	CB17/Icr-Prkdc <sup>scid</sup> /CrIcrlj C3H/HeNCrIcrlj (Sentinels)
	Test period Technique	November 16 - 26
<b>BACTERIA</b>		
<i>Bordetella bronchiseptica</i>	Culture	0/21
<i>Citrobacter rodentium</i>	Culture	0/21
<i>Corynebacterium kutscheri</i>	Culture	0/21
<i>Pasteurella pneumotropica</i>	Culture	0/21
<i>Pseudomonas aeruginosa</i>	Culture	0/21
<i>Salmonella</i> spp.	Culture	0/21
<i>Staphylococcus aureus</i>	Culture	0/21
<i>Streptococcus pneumoniae</i>	Culture	0/21

Tested by Charles River Laboratories Japan, Inc.

Monitoring frequency: Fecal and water samples of isolator-maintained animals are tested monthly.

HEALTH MONITORING REPORT (November 2015)

Printed date: November 28, 2015  
Location: Tsukuba Breeding Center

Species: MICE (Fecal and Water Samples)

Agents	Area	T23 Isolator
	Strains & Stocks	JAX <sup>®</sup> Mice Strain NOD.CB17-Prkdc <sup>scid</sup> /J C3H/HeNCrIcrIj (Sentinels)
	Test period Technique	November 16 - 26
<b>BACTERIA</b>		
<i>Bordetella bronchiseptica</i>	Culture	0/21
<i>Citrobacter rodentium</i>	Culture	0/21
<i>Corynebacterium kutscheri</i>	Culture	0/21
<i>Pasteurella pneumotropica</i>	Culture	0/21
<i>Pseudomonas aeruginosa</i>	Culture	0/21
<i>Salmonella</i> spp.	Culture	0/21
<i>Staphylococcus aureus</i>	Culture	0/21
<i>Streptococcus pneumoniae</i>	Culture	0/21

Tested by Charles River Laboratories Japan, Inc.

Monitoring frequency: Fecal and water samples of isolator-maintained animals are tested monthly.